arXiv:1210.6681v1 [math.GT] 24 Oct 2012

A graphical calculus for tangles on surfaces *

Peter M. Johnson Sóstenes Lins

October 26, 2012

Abstract

We show how the theory of tangles is equivalent to that of well-connected tangles. These are drawn on a surface with boundary, and equivalent via Reidemeister moves of a restricted kind. This reworking of the graphical foundations for link and tangle theory can be expected to have a variety of applications, including ones involving 3-manifolds. It opens the way to new approaches for defining 'facial' state-sum invariants that depend in part on assigning substates to faces of tangle diagrams.

1 Introduction

Our aim is to make a substantial improvement in the foundational part of combinatorial topology currently used to define invariants of knots and links, or of related objects such as tangles on surfaces. The usual assumptions of PL theory, for framed tangles on compact surfaces with boundary (the most general 'knotty' objects of interest to us), apply throughout. We weaken the requirements for defining invariants of tangles on surfaces by making a significant restriction on the allowed Reidemeister moves of type 2 and, to compensate, consider only diagrams that are well-connected, as defined below. To evaluate a new invariant on an arbitrary tangle diagram, the diagram should first be prepared, via unrestricted moves of type 2, so that it becomes well-connected. If that step is omitted, it may still be possible to assign a value to the diagram, without guaranteeing that this defines an invariant.

The present article deals only with the graphical foundations needed to justify the general approach, which does not even require the surfaces to be orientable. A companion article [4] introduces a family of 'facial' state-sum invariants for tangles in oriented surfaces, constructed using the idea introduced here. These arose from a project to abstract and greatly simplify the approach to 3-manifold invariants like those of Witten-Reshetikhin-Turaev, as presented in [6]. The invariants we found resemble those of 2-dimensional TQFT theory, but the decompositions (permitting local calculations that piece together nicely) involve not just surfaces but tangles in surfaces. To calculate with invariants in this family, diagrams need not be prepared to make them well-connected, as it is more efficient to introduce factors that compensate for the effect of dealing directly with faces that have handles or disconnected boundaries. A very simple invariant with interesting properties is constructed explicitly in [4] and studied in detail. It remains to be seen what other approaches can exploit well-connectedness to define new types of invariants.

First some historical background will be sketched. The Reidemeister moves for diagrams of links (including knots) that are generic projections on the plane of links being isotoped in space, were discovered independently by Reidemeister [8] and Alexander and Briggs [1]. They became widely known through Reidemeister's second book on knot theory [9]. The focus is not on a single link or a diagram L but on the class $[L]$ of diagrams equivalent to the given one via appropriate local modifications. Among them we include moves of type 0 (isotopies made to the tangle diagram), type 1 (often restricted or avoided), type 2 (to be restricted below, and subdivided into cases), and type 3. With minor variations, most notably versions where components are oriented, Reidemeister moves usually provide the foundation used to obtain relations whose satisfaction is necessary and sufficient for defining invariants of links and related objects such as framed tangles on surfaces.

An avenue that opens the possibility for obtaining more refined invariants is to calculate only after preparing diagrams in some way. Our main idea involves a new kind of preparation, but first we review some other kinds.

One foundation is provided by presenting links as braids, on the plane or even within 3-manifolds, as in Lambropolou and Rourke [7]. Although we do not do so, ideas about well-connectedness can easily be adapted to apply to braids.

There were several independent discoveries concerning twist and writhe, starting with Călugăreanu [3] in 1959, but these long went unperceived by knot theorists, whose concern was combinatorial topology rather than differential geometry. As late as 1983, Trace [10] showed the irrelevance of the first Reidemeister move, on adding and removing curls. for knot (and hence link) diagrams in the plane. The only Reidemeister moves that matter for defining invariants of oriented links in the plane are those of type 2 and 3, provided the link components are first prepared by adding curls to suitably adjust their writhe and rotation number. In practice, it proved better to ignore the rotation number, to facilitate the study of more general objects such as links on oriented surfaces. The focus shifted to framed objects, where one admits a writhe-preserving form of the first Reidemeister move, also known as a ribbon move. One can always add curls to prepare a diagram so that it represents a link all of whose components are 0-framed. Those who prefer unframed objects, invariant under the full set of Reidemeister moves, can restrict attention to 0-framed diagrams.

---

*2000 Mathematics Subject Classification: 57M25 (primary), 57M27 (secondary)

We are indebted to the Departamento de Matemática, UFPE, Brazil and to the Centro de Informática, UFPE, Brazil for financial support. The second author is also supported by a research grant from CNPq/Brazil, proc. 301233/2009-8.

2 Definitions and results

We will present a new way to prepare diagrams of links or tangles, giving ones that are connected in a strong sense, and will describe which moves are appropriate for equivalence of such diagrams.

Following and adapting usual conventions and definitions for PL objects, tangles (including links) will be drawn on a compact surface S whose boundary ∂S can be regarded as consisting of k disjoint 'circles' left after removing open disks from a surface without boundary. All intersections are transverse, with under/over information recorded at each crossing. Surfaces can be treated abstractly, as in combinatorial topology, but it is also convenient to use informal metric language, for example mentioning two close and almost parallel segments in S. Tangles in S consist of a finite number of components, which are connected curves, each one either closed or with endpoints on the boundary ∂S of S. These are assumed unoriented, merely to avoid listing oriented versions of the basic moves below. We prefer not to mention thickened surfaces until later, so faces of a tangle T on S can be defined to be the connected components of the complement of $T \cup \partial S$ in S.

In S, the boundary of a face relative to T is contained in $T \cup \partial S$, which decomposes naturally into arcs, used in the usual way to codify combinatorial information. This describes how to glue faces at arcs to reconstruct (up to homeomorphism) the tangle in the surface. If one prefers each face to have a boundary in which no two arcs are glued together, consider tangles that have been thickened so that their strands become thin bands on S.

Equivalence of tangles will be defined via Reidemeister moves of types 0 (isotopies of S homotopic to the identity, which also act on T), 2 and 3, giving a version of the regular isotopy classes of Kauffman [5]. Optionally, one could also allow some sort of type 1 move, usually the ribbon move (see for example [6]), which gives the important class of framed tangles. Reidemeister moves are always supposed to modify only the part of the tangle within some open disc.

Tangles (especially links) studied here differ from virtual ones in two ways: it is not permitted to modify S by adding or removing handles on parts of S disjoint from the tangle, and tangles in the same orbit of the mapping class group of S are not necessarily equivalent. The last condition opens the way to defining invariants that in some way use homotopy or homology classes, as in [2]. A useful alternative is to consider the surface S' obtained by removing a point from S, so that a tangle diagram on S' will have a distinguished 'infinite' region. When S' is orientable, it can be obtained from the plane by removing open discs and attaching handles. If S' is nonorientable, there is also a crosscap.

Our main contribution is to remove an obstacle that until now has blocked progress with state-sum invariants that depend in part on assigning states to faces. The problem is a non-local feature of Reidemeister moves of type 2. When two intersecting strands pull apart to form two disjoint strands, three regions (viewed locally) become one. One of these, within the disk, disappears, and the other two fuse. These two may be the same region, globally, but if not the move is said to be *admissible*. We prefer to work with two special kinds of move that are undeniably local: those of types 2E and 2F (E for eight, F for finger), that we now define. Each comes in two forms, with more refined names as shown in the the next figures.



Figure 1: **The 8-moves $2E^+$ and $2E^-$**



Figure 2: **The finger-rotation moves $2F_A$ and $2F_B$**

(2.1) Definition. A tangle T on S is *well-connected* if each face F is an open disk or cylinder whose boundary ∂F in S intersects ∂S in a connected set, which in the second case is a boundary component of S. Two such tangle diagrams on S are *strongly equivalent* if one can be obtained from the other by a sequence of moves of types 0, 2E, 2F and 3.

Local moves will turn out to be sufficient when working with tangles that are well-connected. The illustrations below show part of a well-connected diagram on a surface S, containing a face having one edge (the maximum allowed) in ∂S, that edge being drawn as a small semicircle. The central illustration indicates where admissible moves can be made, while the others can be obtained from their neighbours via moves of type 2F. A barrier is shown where a move between one pair of neighbours is impeded by the presence of an edge in ∂S. The crossing type where strands cross is displayed only when relevant.



Figure 3: **Factoring admissible type 2 moves into 2E and 2F moves**

(2.2) Proposition. *Any move of type 2, applied to a well-connected tangle diagram in the direction that creates a pair of crossings, produces another such diagram. The move can be factored into a move of type 2E followed by a sequence of moves of type 2F.*

Proof. Such a move can be regarded as taking place essentially within a face F, via a finger-creation move where one edge of the face grows within F into a thin finger which then passes under an edge (possibly the one where it originated). The faces adjacent with F need not be different from F, as some pairs of edges around the boundary of F could be identified, but any diagram will at least show faithfully the arrangement of edges around each vertex. All new faces created are topological disks, and the conditions referring to boundaries that intersect ∂S clearly continue to hold. In the typical case shown above, by moving around the outer illustrations, starting with the copy of the central one, any other can be reached. This is done by making a move of type 2E, then moves of type 2F, always within a topological disk. In cases such as the one shown, where a barrier occurs, only one direction to proceed is possible. The only other case of interest is when the region where the move is made (creating crossings) is annular. By definition, this region must have one boundary component in ∂S and the other disjoint from ∂S, and again there is only one direction in which to proceed. ☐

Reidemeister moves of type 2, in the direction that creates a new pair of crossings, can be used to convert tangle diagrams into equivalent ones that are well-connected. It is this process that we call preparation. Sequences of such moves give what will be called finger-creation moves, where a a small segment of T is replaced by a thin finger (almost parallel pair of strands) that expands within S, avoiding ∂S, and passes under T, other fingers, or even earlier parts of the current finger.

(2.3) Proposition. *Let T be a tangle on a surface S such that every connected component of S contains points of T. Then T can be prepared by applying finger-creation moves repeatedly so that the new tangle is well-connected.*

Proof. Given T as above, the procedure is straightforward. One can cut faces until their boundary is connected and intersects ∂S in at most one connected piece. A way to remove handles is shown in the figure below, and it is easy to cut a nonorientable face, say one without handles (a Möbius strip) into two orientable pieces. When a face has a boundary component fully contained in ∂S, finger-creation moves cannot approach that component, and the best that can be done is to isolate the component, creating a face that is topologically a cylinder rather than a disk. ☐



Figure 4: **Cutting a handle by two finger extensions**

(2.4) Theorem. *All prepared diagrams obtained from the same tangle diagram are strongly equivalent.*

Proof. Consider two preparations T_1 and T_2 of the same tangle diagram T. It can be assumed that the small strands of T that were initially expanded are mutually disjoint. Form a new tangle T_3 using both preparations, where any new intersections between T_1 and T_2 are adjusted to be transverse, with the part on T_2 passing under that on T_1. One can then convert both T_1 and T_2 to T_3 via a series of moves of types 2E and 2F between well-connected diagrams, so T_1 and T_2 are strongly equivalent. ☐

This will now be strengthened. We treat only regular isotopy. Analogous results, where further types of moves are admitted, follow immediately.

(2.5) Theorem. *All prepared diagrams of regularly isotopic tangles on S are strongly equivalent.*

Proof. Suppose T' (resp. U') is an arbitrary diagram obtained by preparing T (resp. U) on S, and let $T = T_0, \ldots, T_n = U$ be a sequence of diagrams in which adjacent ones are related by Reidemeister moves of types 0, 2, and 3. We first show how to obtain a sequence $T' = T'_0, \ldots, T'_n$ of well-connected tangle diagrams, where each T'_i is a diagram prepared from T_i by using new strands that always pass under the ones of T_i, and consecutive diagrams T'_i, T'_{i+1} are related by a series of restricted Reidemeister moves. After the given initial T', each T'_{i+1} in the sequence is obtained from T'_i by carrying out moves related to the one between T_i and T_{i+1}, after moving out of the way any new strands that obstruct the move (passing under the others). All this can clearly be done using only moves of types 0, 2E, 2F, and 3. Finally, note that T'_n and U' are preparations of the same tangle U. Thus T' and U' are strongly equivalent. $\square$

As an obvious reformulation of the above ideas, we have:

(2.6) Corollary. *Each regular isotopy class* $[T]$ *of tangles on* S *determines, via preparations, a unique strong equivalence class* $\langle T' \rangle$ *of tangles on* S*. The members of the latter class are the well-connected tangles contained in the first.*

The class $\langle T' \rangle$ can be regarded as a graphical invariant of the original regular isotopy class $[T]$. This is a more promising starting point for methods that construct various families of algebraically-defined invariants, as only certain kinds of Reidemeister moves need be respected. In a companion article [4], the first in a planned series, we introduce invariants of this form.

References

[1] J.W. Alexander and G.B. Briggs. On types of knotted curves. *Annals of Mathematics*, 28(2):562–586, 1927.

[2] M. Barone and M. Ferri. Homological Farrell invariants for embedded graphs. *Atti Sem. Mat. Fis. Univ. Modena*, 49:217–227, 2001.

[3] G. Călugăreanu. L'intégral de gauss et l'analyse des noeuds tridimensionnels. *Rev. Math. Pures Appl.*, 4:5–20, 1959.

[4] P. Johnson and S. Lins. A state sum invariant of tangles in surfaces. *In preparation*, 2012.

[5] L.H. Kauffman. An invariant of regular isotopy. *Trans. Amer. Math. Soc*, 318(2), 1990.

[6] L.H. Kauffman and S. Lins. Temperley-Lieb Recoupling Theory and Invariants of 3-manifolds. *Annals of Mathematical Studies, Princeton University Press*, 134:1–296, 1994.

[7] S. Lambropolou and P. C. Rourke. Markov's theorem in 3-manifolds. *Topology and its Applications*, 78:95–122, 1997.

[8] K. Reidemeister. Elementare begründung der knotentheorie. *Abh. Math. Sem. Univ. Hamburg*, 5:24–32, 1926.

[9] K. Reidemeister. *Einführung in die kombinatorische Topologie.* Springer, 1932.

[10] B. Trace. On the Reidemeister moves of a classical knot. *Proc. Amer. Math. Soc.*, 89:722–724, 1983.

Peter M. Johnson
Departamento de Matemática, UFPE
Recife–PE
Brazil
peterj@dmat.ufpe.br

Sóstenes Lins
Centro de Informática, UFPE
Recife–PE
Brazil
sostenes@cin.ufpe.br